



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME

UNITED STATES DEPARTMENT OF COMMERCE
United States Patent and Trademark Office

November 22, 2004

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A
FILING DATE.

APPLICATION NUMBER: 60/513,779
FILING DATE: October 23, 2003
RELATED PCT APPLICATION NUMBER: PCT/US04/34975

Certified by



Jon W Dudas

Acting Under Secretary of Commerce
for Intellectual Property
and Acting Director of the U.S.
Patent and Trademark Office



BEST AVAILABLE COPY

102303

16085 U.S. PTO

PTO/SB/16 (08-03)

Approved for use through 07/31/2008. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No.

EV 0103 20211 US

2008 U.S. PTO
60/513767

102303

INVENTOR(S)					
Given Name (first and middle (if any))		Family Name or Surname		Residence (City and either State or Foreign Country)	
Sidney		LIANG		Olivette, Missouri, USA	
Additional inventors are being named on the <u>0</u> separately numbered sheets attached hereto					
TITLE OF THE INVENTION (500 characters max)					
IMPROVED PROCESS FOR PREPARING BENZHYDRYLTHIOACETAMIDE					
Direct all correspondence to: CORRESPONDENCE ADDRESS					
<input checked="" type="checkbox"/> Customer Number: 24289					
OR					
<input type="checkbox"/> Firm or Individual Name					
Address					
Address					
City		State		Zip	
Country		Telephone		Fax	
ENCLOSED APPLICATION PARTS (check all that apply)					
<input checked="" type="checkbox"/> Specification Number of Pages <u>14</u> <input type="checkbox"/> CD(s), Number _____					
<input type="checkbox"/> Drawing(s) Number of Sheets _____					
<input type="checkbox"/> Application Date Sheet. See 37 CFR 1.76					
<input type="checkbox"/> Other (specify) _____					
METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT					
<input type="checkbox"/> Applicant claims small entity status. See 37 CFR 1.27.					
<input type="checkbox"/> A check or money order is enclosed to cover the filing fees.					
<input checked="" type="checkbox"/> The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: <u>13-1160</u>					
<input type="checkbox"/> Payment by credit card. Form PTO-2038 is attached.					
<div style="float: right; border: 1px solid black; padding: 5px; text-align: center;"> FILING FEE Amount (\$) \$160.00 </div>					
The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.					
<input checked="" type="checkbox"/> No.					
<input type="checkbox"/> Yes, the name of the U.S. Government agency and the Government contract number are: _____					

[Page 1 of 2]

Respectfully submitted:

SIGNATURE

TYPED or PRINTED NAME Jeffrey S. BooneTELEPHONE 314-654-8955Date 10/22/03REGISTRATION NO. 29,284

(If appropriate)

Docket Number: 1661 P US

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

IMPROVED PROCESS FOR PREPARING BENZHYDRYLTHIOACETAMIDE

[0001] This invention relates to an improved process for preparing modafinil wherein the intermediate, benzhydrylthioacetate is prepared in a reaction medium comprising an organic solvent and water having dissolved therein a basic salt. The acetate thus produced is amidated and then oxidized to provide benzhydrylsulphonylacetamide, commonly known as modafinil.

Background Of The Invention

[0002] Lafon disclosed modafinil and other similar compounds in U.S patent 4,177,290 as having pharmaceutical activity on the central nervous system. In a typical prior art process, benzhydrylthioacetic acid is halogenated with thionyl chloride. The chloride is then converted to the amide in methylene chloride with ammonia. The amide is then oxidized with hydrogen peroxide to provide benzhydrylsulphonylacetamide. Other derivatives of modafinil as well as methods of preparation and purification are disclosed in U. S. patent 4,127,722. However, the amide appears to be the compound of choice among the many derivatives now known.

[0003] Interest in the Lafon compounds has increased in recent years because these compounds have been discovered to have beneficial effects in the treatment of a wide variety of diseases in mammals including humans. Although first noted as a treatment for narcolepsy, more recent patents and technical publications have listed such compounds as beneficial in the treatment of Parkinson's disease, urinary incontinence, Alzheimer's disorder, ischemia and stroke. As the use of these compounds increased so has the demand for greater volumes while maintaining the highest state of purity and also avoiding process chemicals of high environmental risk.

[0004] Lafon also disclosed the use of an intermediate, a racemic mixture of benzhydrylsulphonylthioacetic acid to prepare an isomer of modafinil in U.S. Patent 4,927,855, referring

to French patent 2,326,181B. The racemic mixture of the acid was reacted with the (-)- α -methylbenzylamine to provide the (-)-benzhydrylsulphinylacetate of (-)-methylbenzylamine which is acidified with concentrated hydrochloric acid to provide (-)-benzhydrylsulphinylacetic acid. The levo- acid thus produced was then reacted with methylsulfate in water and sodium bicarbonate to provide methyl(-)-benzhydrylsulphinylacetate. The acetate was then amidated with ammonia gas to provide (-)-benzhydrylsulphinylacetamide.

[0005] Numerous substituted thioacetamides are disclosed in US 6,492,396 to Bacon et al. In one synthesis scheme benzhydrol is converted to a benzhydrylthiol by reaction with thiourea that is then converted by hydrolysis to a thiouronium moiety. Subsequently, the thiouronium is converted to an acid with chloroacetic acid. The benzhydrylthioacetic acid is treated in various ways depending upon the desired derivative. To prepare the amide the acid is reacted with ammonia or an appropriate amine in an organic solvent such as tetrahydrofuran or methylene chloride. Other thioacetamide derivatives are obtained by employing N-methylmorpholine and a thioacetic acid in dimethylformamide (DMF).

[0006] A procedure for the preparation of an acetamide intermediate for the production of modafinil is disclosed in published US application 2002/0183552. According to this application a three-step procedure for preparing modafinil is disclosed starting with benzhydrol (diphenylmethanol) that is employed to prepare the benzhydrylthiocarboxamidine salt by reaction with thiourea in hydrogen bromide. The bromide salt is then reacted with chloroacetamide in aqueous sodium hydroxide to produce diphenylmethylthioacetamide. The acetamide may then be oxidized by conventional means to produce modafinil. Typically, the oxidation is provided by a reaction with hydrogen peroxide in glacial acetic acid.

[0007] Because of the growing demand for large quantities of modafinil in a highly pure state there is needed a process for preparing the product efficiently without need for undesirable starting

materials or by-products. In particular the production of the acetamide intermediate is particularly in need of improvement although given some degree of attention in the art as indicated above.

Summary Of The Invention


[0008] In accordance with this invention there is provided a novel process for the preparation of benzhydrylthioacetate by the reaction of a haloacetate with the reaction product of thiourea and benzhydrol. The improved process is conducted in a solvent comprising water, having dissolved therein a basic salt and a water miscible organic solvent. The resulting benzhydrylthioacetate is provided in high yield and purity and can be amidated and oxidized to provide the pharmaceutical modafinil.

[0009] The process of this invention provides improved yield and purity over known processes. The water miscible organic solvent provides the dual function of providing a solvent for the starting organic material while also providing a convenient reaction medium allowing low temperature reactions to take place.


Detailed Description Of The Invention

[0010] The pharmaceutical modafinil is conveniently prepared in three steps including the novel step of this invention wherein 2-chloroacetate, a water-soluble organic solvent and a basic salt is employed. First, benzhydrol 1 is reacted with thiourea 2 to provide a reaction product that is sometimes termed the benzhydrylthiocarboxamidine salt 3. This reaction is carried out in water in the presence of hydrogen halide, shown below as the bromide at a temperature of about 80°C for a period of from about 1 to about 2 hours. However, it has been found that the chloride is preferred in this first step because there is produced less by product and higher efficiency. A solid


benzhydrylthiocarboxamide halide salt precipitates. The reaction may be described structurally as



[0011] In the second step, the benzhydrylthiocarboxamide bromide **3** salt is washed with water then placed in a reaction vessel together with methyl 2-chloroacetate **4**, a basic salt, an organic, water soluble solvent such as methanol and water. This mixture is stirred to dissolve the bromide salt and to allow it to react at or near room temperature for 18 hours to provide methyl benzhydrylthioacetate. Although the reaction is described with respect to the bromide salt, any other suitable salt may be employed. After formation of the acetate, it can be converted to the amide with ammonia gas. This is accomplished by passing ammonia gas through the reaction mixture for about 1 hour followed by stirring for an additional 5 hours. When the reaction is completed the reaction mixture is diluted with water and a solid precipitate is separated. When washed with water, benzhydrylthioamide **5** is obtained in high yield and purity. The reaction may be described structurally as follows:



[0012] In the third step of the procedure to obtain modafinil in accordance with this invention, benzhydrylthioacetamide **5** is dissolved in acetic acid and then hydrogen peroxide is slowly charged to the solution while cooling to control the exothermic reaction. Typically the temperature is maintained below 22°C. to prevent undesired side reactions. After the reaction is completed, diluting the reaction mixture with water and separating the precipitate to obtain a crude racemic benzhydrylsulphinylacetamide **6** (modafinil) isolates the product. The crude product is typically refined by recrystallization in a solvent or a mixture of solvents including chloroform to obtain highly pure pharmaceutical grade modafinil in about 70% overall yield. The reaction may be described structurally as follows:



[0013] The term "alkali metal" encompasses lithium, sodium, potassium, cesium and rubidium; and the term "alkaline earth metal" includes beryllium, magnesium, calcium, strontium and barium. Typically, the preferred alkali metals are sodium and potassium while the preferred alkaline earth metal salts are calcium and magnesium.

[0014] Any suitable water miscible organic solvents can be employed in the process of this invention. The water miscible organic solvent must have some degree of solvating power with respect to the organic starting material benzhydrylthiocboxamidine salt. Typically, such solvents include acetone and lower alkanols such as methanol, ethanol, butanol, sec-butyl alcohol and tert-butyl alcohol. However, it has been found that dimethylformamide(DMF) is particularly useful as well as other water miscible amides.

[0015] The process of this invention, employing a water miscible organic solvent/water reaction medium containing a basic salt, can be run at room temperature and produces isolated yields in the range of about 97% (based upon HPLC analysis) with the amount of each impurities at less than 3%. Such results represent significant yield improvements as well as a more efficient process compared to all other known methods. This process eliminates a process step and the use of thionyl chloride and benzene in comparison to the '290 patent. The process of this invention also eliminates the use of high temperatures, corrosive solutions, and byproducts, while still producing higher yields with very low impurities compared to the prior art processes such as those that employ sodium hydroxide with haloacetamide in the above described published application.

[0016] Any number of basic salts can be employed in the process of this invention. Salts may be employed that may only be slightly soluble in water. However, the presence of the salt in the reaction medium remains effective to promote the reaction in the particulate form. Thus the term "water soluble" as employed herein includes material only slightly soluble in water. In particular, it is preferred to employ ammonium, alkali metal salts or alkaline earth metal salts. In particular, the sodium salt is preferred and the potassium salt is even more preferred. The anion of the salt is typically a sulfate, sulfide, phosphate, bicarbonate, nitrate, phosphonate, phosphinate and preferably a carbonate. Typical salts included in the above description of water soluble salts are sodium sulfate, calcium sulfate, magnesium sulfate, sodium sulfide, magnesium sulfide, calcium sulfide, sodium phosphate, magnesium phosphate, calcium phosphate, potassium phosphate, sodium bicarbonate, calcium bicarbonate, magnesium bicarbonate, sodium nitrate, calcium nitrate, magnesium nitrate, sodium phosphonate, potassium phosphonate, magnesium phosphonate, calcium phosphonate, sodium phosphinate, potassium phosphinate, calcium phosphinate, magnesium phosphinate, potassium sulfate, potassium sulfide, potassium bicarbonate, potassium nitrate, potassium tripolyphosphate, sodium tripolyphosphate, sodium thiophosphate, potassium citrate, tetrapotassium pyrophosphate, ammonium phosphate, ammonium chloride, ammonium sulfate, ammonium bicarbonate,

ammonium phosphinate, ammonium phosphonate and the like. Potassium carbonate is preferred because it is readily soluble in water.

[0017] The process of this invention has been found to be particularly advantageous over the prior art sodium hydroxide process. Said prior art process requires higher temperatures to help increase the yields since lower temperatures tend to produce significant amounts of impurities. A milder basic solution was thought to be sufficient enough to promote the reaction and at lower temperature conditions. Water and potassium carbonate were initially used, but byproducts were still produced. In a preferred embodiment, a mixture of methanol and potassium carbonate solution was employed to help dissolve the starting reagent and promote a more stable pH during the reaction. The amount of basic salt, by weight of the total the starting reagent benzhydrylthiocarboxamidine bromide salt employed in the process of this invention, is typically in the range of from about 41% to about 200% and preferably in the range of from about 82% to about 110% and most preferably in the range of from about 95% to about 105%.

[0018] In another aspect of this invention is the clean up of the final product, modafinil. It has been found that mixing it with chloroform best purified the final product. The preferred method is to then refluxing the mixture for a short period of time. The refluxed mixture is then cooled to a relatively low temperature, filtered and washed to provide a highly pure modafinil product. The use of methanol and/or methanol/water solvent as disclosed in the original '290 patent to purify the modafinil was found to be inadequate or inefficient in obtaining pharmaceutically pure modafinil. Similar alcohol solvents such as ethanol and propanol also gave similar results wherein several impurities were significantly greater than 0.1%. Modafinil was only mildly soluble, at best, in alcoholic solvents even at reflux temperatures. Many impurities were also very insoluble in alcoholic solvents and were thus retained in modafinil. In many cases 1g of crude modafinil required a minimum of 8ml of methanol to be completely dissolved at reflux temperature. When filtered at room temperature, many impurities were only moderately reduced.

[0019] It also been found that excellent purification of modafinil can be achieved by mixing the crude modafinil product with a halo-organic solvent such as dichloromethane, dichloroethane and preferably chloroform. Chloroform was initially believed to be a better solvent for the clean-up procedure because it was slightly acidic in comparison to alcohols. Surprisingly, modafinil was also extremely insoluble in chloroform, but fortunately, the impurities were very soluble in chloroform. Major impurities in the crude modafinil product include the modafinil acid, modafinil sulfone acid, modafinil sulfone and unreacted starting material, benzhydrylthioacetamide. Crude modafinil was initially mixed with chloroform in 1g crude to 4 ml chloroform. It has been found that the major impurities are substantially removed by the chloroform washing.

[0020] In a preferred embodiment a low boiling aliphatic solvent, preferably heptane, can be added in a ratio of about 2ml of solvent to about 1g crude to help reduce the viscosity of the slurry. To address the problem of the viscous slurry, heptane can be charged first to the crude. Chloroform is then added to the stirred mixture slowly. A mild slurry, results. Heating the slurry to reflux further alleviates the viscosity to a simple solid/liquid mixture even when it was cooled to 5°C allowing easy filtering with high yields. The slurry would become less viscous only upon heating to reflux for about 30 minutes. Even though the modafinil never completely dissolves in the solvent mixture, the cleaning procedure is effective. The solvent mixture is then cooled to about 5°C allowing the modafinil to precipitate fully after which it can be filtered to obtain 92% to 97% recovery yield with ~99.8% purity by weight (HPLC analysis), respectively. In this preferred embodiment any suitable low boiling aliphatic solvent such as pentane, hexane, heptane, or octane may be employed.

[0021] The following examples are intended to illustrate the present invention and are not to limit the claims in any manner. All of the percentages are by weight unless otherwise indicated.

Preferred Embodiments Of The Invention

[0022] EXAMPLE 1

Preparation of benzhydrylthiocarboxamidine salt.

[0023] To a reaction vessel there was charged 82.63g of thiourea, 150 ml of HBr (48%) and 200 ml of water. Then, 100g of benzhydrol were charged to the mixture in the reaction vessel. The reaction mixture was then refluxed at 90°C for 5 hr. The reaction mixture was then cooled to room temperature and 100 ml of additional water were added to the mixture followed by filtration. The crude solid residue was rinsed with 75 ml and air dried to provide approximately 90%(molar) yield at greater than 95% purity.

[0024] EXAMPLE 2

Preparation of benzhydrylthioacetamide

[0025] To a 1 liter 3-neck round bottom flask there were added 86.5 g of benzhydrylthiocarboxamidine bromide from Example 1, 35.3 g of methyl chloroacetate, 93.5 g of potassium carbonate and 300 ml of methanol. The reaction mixture was maintained at room temperature in a water bath and continued stirring at room temperature for 24 hours. The resulting solution exhibited a slight, pink color. The reaction mixture was then cooled to 15°C and ammonia gas was bubbled through the reaction for 1 hour at a high rate. The reaction was diluted with 600 ml of water. A precipitate formed and the reaction mixture was filtered to obtain 58g of product. Analysis by HPLC indicated the presence of benzhydrylthioacetamide in 89% purity.

[0026] EXAMPLE 3

Preparation of benzhydrylsulphinylacetamide

[0027] To a 500ml three-necked round bottom flask was charged 50 grams of benzhydrylthioacetamide and 100 ml of acetic acid. The mixture was stirred until all solids were dissolved and then the reaction mixture was cooled to 15°C. There were then slowly added to the reaction mixture 25 ml of hydrogen peroxide solution (30%) in step-wise fashion (5-10ml portions)

while maintaining the temperature of the reaction mixture below 20°C. The reaction mixture was then stirred at room temperature (20°C) or until the amide disappeared. There was then added 500 ml of water to the reaction mixture thereby precipitating the product. The reaction mixture was cooled to 15°C and filtered. The crude solid product was then rinsed with 50 ml of water. The product was then purified by combining it with heptane (3.5ml/1g crude) and then chloroform (7ml/1g of crude). The mixture was refluxed for 30 min. at a temperature of 70-75°C. The solution was slowly cooled to 10°C with stirring and the solid precipitate then filtered and oven dried. The yield was 85%(molar) and the purity was 99.8%.

[0028] Although the invention has been described in terms of specific embodiments which are set forth in considerable detail, it should be understood that this description is by way of illustration only and that the invention is not necessarily limited thereto since alternative embodiments and operating techniques will become apparent to those skilled in the art in view of the disclosure. Accordingly, modifications are contemplated which can be made without departing from the spirit of the described invention.

What is Claimed is:

1. A process for the preparing of benzhydrylthioacetate comprising reacting benzhydrylthiocarboxamidine salt with haloacetate in a reaction medium comprising water, a water miscible organic solvent and a water soluble basic salt.
2. The process of Claim 1 wherein the basic salt is selected from the group consisting of alkali metal and alkaline earth metal sulfates, sulfides, phosphates, carbonates, bicarbonates, nitrates, phosphonates and phosphinates.
3. The process of Claim 1 wherein the haloacetate is 2-chloroacetate.
4. The process of Claim 2 wherein the basic salt is a potassium salt.
5. The process of Claim 4 wherein the basic salt is a potassium carbonate salt.
6. The process of Claim 1 wherein the basic salt is present in a weight ratio to benzhydrylthiocarboxamidine salt of from about 41 to about 200.
7. The process of Claim 6 wherein the basic salt is present in the weight ratio of from about 82 to about 110.
8. The process of Claim 6 wherein the basic salt is present in the weight ratio of about 105.
9. The process of Claim 1 wherein the water miscible organic solvent is selected from the group consisting of lower alkanols, acetone and dimethylformamide.
10. The process of Claim 12 wherein the lower alkanol is selected from the group consisting of methanol, ethanol, butanol, sec-butyl alcohol, tert-butyl alcohol, and acetone.
11. A process for preparing modafinil comprising the following steps:
 - a) reacting benzhydrol with thiourea in the presence of hydrogen halide to provide benzhydrylthiocarboxamidinehalide;
 - b) reacting haloacetate with the product of step a) to provide benzhydrylthioacetate and without isolation, amidating the acetate to obtain benzhydrylthiolacetamide and,

c) oxidizing the product of step b) to obtain benzhydrylsulphinylacetamide.

12. The process of Claim 14 wherein the basic salt is selected from the group consisting of alkali metal and alkaline earth metal sulfates, sulfides, phosphates, carbonates, bicarbonates, nitrates, phosphonates and phosphinates.
13. The process of Claim 11 wherein the haloacetate is 2-chloroacetate.
14. The process of Claim 12 wherein the basic salt is a potassium salt.
15. The process of Claim 14 wherein the basic salt is a potassium carbonate salt.
16. The process of Claim 12 wherein the basic salt is present in a weight ratio to benzhydrylthiocarboxamidine salt of from about 41 to about 200.
17. The process of Claim 16 wherein the basic salt is present in the weight ratio of from about 82 to about 110.
18. The process of Claim 17 wherein the basic salt is present in the weight ratio of about 105.
19. The process of Claim 11 wherein the water miscible organic solvent is selected from the group consisting of lower alkanols, acetone and dimethylformamide.
20. A process of Claim 11 further including the step of purifying the product of step c0 which comprises contacting the crude product with a halo-organic solvent and then separating the modafinil from the solvent.
21. The process of Claim 20 wherein the temperature of the mixture of the product and halo-organic solvent is raised to a reflux temperature.
22. The process of Claim 21 wherein the reflux temperature is maintained for about 30 minutes.
23. The process of Claim 20 wherein the halo-organic solvent is selected from the group consisting of chloroform, dichloromethane, and dichloroethane.
24. The process of Claim 20 further including the step of adding an aliphatic solvent to the mixture.

25. The process of Claim 24 wherein an aliphatic solvent is added to product prior to contacting the product with the halo-organic solvent and the temperature of the mixture is raised to the reflux temperature.
26. The process of Claim 25 wherein the aliphatic solvent is selected from the group consisting of pentane, hexane, heptane and octane.
27. The process of Claim 24 wherein the halo-organic solvent is chloroform and the aliphatic solvent is heptane.

IMPROVED PROCESS FOR PREPARING BENZHYDRYLTHIOACETAMIDE

Abstract Of The Invention

[0029] The present invention is directed to an improved process for preparing modafinil wherein benzhydrylthioacetate is prepared in high yield and purity by the reaction of a haloacetate with the reaction product of thiourea and benzhydrol in aqueous solution. The reaction employing the haloacetate is conducted in a solvent comprising water, an organic solvent such as methanol having dissolved therein a basic salt such as potassium carbonate. The resulting benzhydrylthioacetate can be amidated and then oxidized to provide the pharmaceutical modafinil in high yield and purity.

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US04/034975

International filing date: 22 October 2004 (22.10.2004)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/513,779
Filing date: 23 October 2003 (23.10.2003)

Date of receipt at the International Bureau: 01 December 2004 (01.12.2004)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse